

PCR and RT-PCR in the Diagnosis of Laboratory Animal Infections and in Health Monitoring

Susan R Compton*

Molecular diagnostics (PCR and RT-PCR) have become commonplace in laboratory animal research and diagnostics, augmenting or replacing serological and microbiologic methods. This overview will discuss the uses of molecular diagnostics in the diagnosis of pathogenic infections of laboratory animals and in monitoring the microbial status of laboratory animals and their environment. The article will focus primarily on laboratory rodents, although PCR can be used on samples from any laboratory animal species.

Abbreviations: IVC, individually ventilated cage; MALDI-TOF, Matrix-Assisted Laser Desorption/Ionization-Time of Flight Mass Spectrometry; MAP, mouse antibody production; MHV, mouse hepatitis virus; MNV, murine norovirus; MPV, mouse parvovirus

DOI: 10.30802/AALAS-JAALAS-20-000008

The use of molecular methods (PCR and RT-PCR) to diagnose bacterial, viral, fungal and parasitic infections in laboratory animals has become common, frequently replacing or augmenting serological, microscopic and culture-based methods.²⁴ Because the majority of laboratory animals are rodents, this review will focus on the detection of infectious agents in rodents and their environment, but PCR and RT-PCR testing of many other commonly used laboratory species such as rabbits, dogs, cats, ferrets, pigs, fish, frogs, and nonhuman primates are commercially available. Recommendations on which infectious agents to test for and exclude from mouse and rat colonies, and the frequency with which the testing should be performed, have been developed by FELASA and AALAS.^{44,117} PCR is used to monitor for over 60 rodent infectious agents (Figure 1).^{18,44,71} FELASA health monitoring guidelines for hamsters, guinea pigs, rabbits, and nonhuman primates have also been published.^{4,44}

PCR or RT-PCR can overcome many of the limitations of other methods of pathogen detection (Figure 2) and therefore represent a refinement of health monitoring in the vivarium. Briefly, PCR-based diagnostic methods amplify the genome of viruses present in, or released from, infected cells or the DNA of bacteria, fungi, or parasites.^{24,53} Three advantages of PCR are its speed, sensitivity, and specificity; unlike serologic or microbiologic techniques, molecular diagnostics can amplify a specific region of a gene from a few copies to trillions in a few hours. However, the high sensitivity of PCR can lead to false positives if samples or reagents are contaminated with nucleic acids from the infectious agent (positive controls) or products from recent PCR assays.⁸⁰ Because RNA is prone to degradation by RNases, RT-PCR can be prone to false negatives if care is not used to prevent RNA degradation during sample collection and RNA extraction. PCR is versatile and can be used on tissue samples or excreta from any species of animal or from samples taken from the laboratory animal environment. Collecting samples for PCR testing requires knowledge of which

tissues become infected with the agent of interest and where the pathogen is shed. The DNA primers selected for PCR determine its specificity.^{24,53} Primers are designed based on available sequence data and can either amplify a single species of an agent (for example, *Mycoplasma pulmonis*) or amplify several related agents (all *Mycoplasma* species). For rapidly evolving agents such as RNA viruses, a nucleotide change in the primer region can lead to a false-negative result in that the agent is present in the sample, but the targeted gene is not amplified.

This review will not discuss methods for nucleic acid extraction and amplification, or their limitations, as these methods have been published previously^{53,106} and are provided with commercial nucleic acid extraction and amplification kits.

Viral infections

The clinical signs and/or histopathology produced by some viruses are specific to a single virus. For example, the characteristic dermal lesions of Ectromelia virus,⁴⁸ the chromodacryorrhea and sneezing caused by rat coronavirus,¹⁰ the enterocyte syncytia in young mice caused by mouse hepatitis virus (MHV)⁵ or the acidophilic intranuclear inclusions in the salivary glands cause by mouse and rat cytomegalovirus¹⁵ can all be used to make a presumptive diagnosis. However, viruses that cause severe distinctive lesions essentially have been eliminated from laboratory rodent colonies, resulting in focus on viruses that cause little or no pathology or morbidity. Therefore, diagnosis of viral infections in rodents using serology has become the most common method of diagnosis.^{24,55,129}

Serologic methods detect antibodies specific for the infectious agent in blood collected from an animal. Serology is an indirect method for detecting infection; therefore, knowing the tissue that harbors the infection is not necessary. Because species-specific secondary antibodies are used in most serology tests, they are limited to those species for which commercial sources of species-specific antibodies are available.

Virus-specific antibodies are first detected in immunocompetent mice 5 to 7 d after infection and peak levels of antibodies are detected 10 to 20 d after infection. Because antibodies may persist for months, a seropositive animal may not be acutely

Infectious Agent	Agent Type	Monitored in Mice	Monitored in Rats
Boone cardiovirus	Virus		X
<i>Bordetella bronchoseptica</i>	Bacteria	X	X
<i>Bordetella hinzii</i>	Bacteria	X	
<i>Bordetella pseudohinzii</i>	Bacteria	X	
Campylobacter	Bacteria	X	X
Chilomastix	Parasite	X	X
<i>Citrobacter rodentium</i>	Bacteria	X	
<i>Clostridium piliforme</i>	Bacteria	X	X
<i>Corynebacterium bovis</i>	Bacteria	X	
<i>Corynebacterium kutscheri</i>	Bacteria	X	X
Cryptosporidium	Parasite	X	X
Dermatophytes	Fungi	X	X
Ectromelia virus	Virus	X	
<i>Encephalitozoon cuniculi</i>	Fungi		X
<i>Entamoeba muris</i>	Parasite	X	X
<i>Filobacterium rodentium</i>	Bacteria	X	X
Fur mites (Myobia/Myocoptes/Radfordia)	Parasite	X	X
<i>Giardia muris</i>	Parasite	X	X
H-1 parvovirus	Virus		X
Hantaviruses (Hantaan/Seoul)	Virus	X	X
Helicobacter	Bacteria	X	X
K virus	Virus	X	
<i>Klebsiella oxytoca</i>	Bacteria	X	X
<i>Klebsiella pneumoniae</i>	Bacteria	X	X
Lactate dehydrogenase-elevating virus	Virus	X	
Lymphocytic choriomeningitis virus	Virus	X	
Minute virus of mice	Virus	X	
Mouse hepatitis virus	Virus	X	
Mouse kidney parvovirus	Virus	X	
Mouse parvovirus	Virus	X	
Murine polyomavirus	Virus	X	
Mouse thymic virus	Virus	X	
Murine adenoviruses 1 and 2	Virus	X	X
Murine chapparvovirus	Virus	X	
Murine cytomegalovirus	Virus	X	
Murine norovirus	Virus	X	
Murine rotavirus	Virus	X	
<i>Mycoplasma pulmonis</i>	Bacteria	X	X
Other mites (Demodex/Ornithonyssus)	Parasite	X	X
Pinworms (Aspicularis/Syphacia)	Parasite	X	X
<i>Pneumocystis murina</i>	Fungi	X	X
Pneumonia virus of mice	Virus	X	X
<i>Proteus mirabilis</i>	Bacteria	X	X
<i>Pseudomonas aeruginosa</i>	Bacteria	X	X
Rat coronavirus	Virus		X
Rat minute virus	Virus		X
Rat parvovirus	Virus		X
Rat polyomavirus 2	Virus		X
Rat virus	Virus		X
Reovirus	Virus	X	X
<i>Rodentibacter pneumotropicus and heylii</i>	Bacteria	X	X
Salmonella	Bacteria	X	X
Sendai virus	Virus	X	X
<i>Spiroplasma muris</i>	Parasite	X	X
<i>Staphylococcus aureus</i>	Bacteria	X	X
<i>Staphylococcus xylosum</i>	Bacteria	X	X
<i>Streptobacillus moniliformis</i>	Bacteria	X	X
<i>Streptococcus</i> , β hemolytic	Bacteria	X	X
<i>Streptococcus pneumoniae</i>	Bacteria	X	X
Theilovirus	Virus	X	X
<i>Trichomonas muris</i>	Parasite	X	X

Data derived from FELASA, Charles River and IDEXX recommended panels [references 18, 44 and 71]

Figure 1. Infectious agents recommended for monitoring in mice and rats.

infected and the infectious agent may have already been cleared. The presence of viral antibodies in a weanling mouse or rat may represent transfer of maternal antibodies from a previously

infected dam, rather than infection of the weanling. Serologic testing is not appropriate for immunocompromised or neonatal mice that do not mount an adequate antibody response. While

	Pathogen types detected	Used with environmental samples	Pooled samples	Used in wide range of animal species	Time to diagnosis	Pathogen identification
PCR/RT-PCR	Viruses, Bacteria, Fungi, Parasites	Yes	Yes	Yes	Hours	Pathogen-specific primers and sequencing
Histopathology	Viruses, Bacteria, Fungi, Parasites	No	No	Yes	Days	Visualization of lesions in stained tissues
Growth in/on culture media	Bacteria and Fungi	Yes	Yes	Yes	Days to weeks	Growth conditions, stains, biochemical testing of isolates
Growth in cultured cells	Primarily Viruses	No	No	Restricted by culture cell type availability	Days to weeks	General identification based on cell type and cytopathic effects observed
Serology	Primarily Viruses and Selected Bacteria and Parasites	No	Yes	Restricted by species specific secondary antibody availability	Hours (if infection is at least 5 days old)	Pathogen-specific antibodies
Electron microscopy	Primarily Viruses	No	No	Yes	Days	Visualization of pathogen
Light microscopy	Parasites	No	Yes	Yes	Hours	Visualization of pathogen

Figure 2. Laboratory animal infectious agent detection methods.

serological responses can be quantified, a stronger antibody response does not correlate with the presence of higher viral loads in the animal. Soiled-bedding sentinel serology is used in many laboratory rodent facilities for health monitoring, as sera from a single rodent can be used to determine whether an infection occurred in any of the cages of experimental mice or rats from which soiled bedding was collected. A limitation of soiled bedding sentinels is that viruses that are not transmitted via the fecal-oral route, such as Sendai virus, lymphocytic choriomeningitis virus, and rat coronavirus, will not be detected effectively.^{36,72,81}

The growth of viruses in cultured cells can be used to quantify the amount of virus produced in an animal.¹²⁸ Detection of viral infections by culture is best attempted early in the infection, prior to the production of an immune response against the virus. However, culture-based methods of viral detection are limited by the fact that not all viruses can be grown in cultured cells, and few laboratory animal laboratories are equipped to cultivate viruses. Furthermore, many viruses do not produce visible cytopathic effects in cultured cells or must be passaged multiple times in cultured cells before cytopathic effects are evident, making it difficult to determine if an infection has occurred. Moreover, if a sample is being screened for multiple viruses, several different cell types may be needed to culture all viral species that are present. Once a virus has been isolated from cultured cells, other methods such as serology or PCR are used to speciate the virus.

PCR of viral nucleic acids extracted from infected tissue can detect the presence of a virus far more rapidly than culture-based methods. Culture-based methods of diagnosis only detect infectious forms of the virus, and days or weeks of incubation may be necessary before virus is detectable.¹²⁸ In contrast, PCR detects infectious viruses, noninfectious forms of the virus and viral transcripts present in infected cells. For PCR detection, knowing what tissues are infected by the virus is crucial so that the correct sites are sampled.⁵³ Quantitative PCR (qPCR or qRT-PCR) can be used to measure the amount of viral nucleic acids present in the sample. While the number of viral genomes and the number of infectious virus particles are usually not equal, PCR can be used to compare the viral titers in viral stocks or tissues. PCR is useful for determining the relative titer of viruses that cannot be cultured in vitro (for example, lactate dehydrogenase-elevating virus and murine astrovirus).^{22,109,133} During an outbreak, PCR followed by sequencing of the amplified product can be used to determine whether the strain of virus is the same in 2 groups of animals (for example, established colony animals compared with newly imported animals).²⁵ The ability of PCR to detect viral infections before seroconversion occurs can be used to determine which animals should be removed from the animal room to decrease the risk of viral transmission within a

laboratory animal population. For some viruses, such as mouse parvovirus, not all strains of mice are equally susceptible to infection. The duration of infection and amount of virus shed is also variable among strains,^{85,87} so testing multiple strains of mice may be necessary to detect the presence of an infectious agent. Thus, multiple rounds of testing are recommended before considering a facility free of an infectious agent.⁸⁵

PCR can be used to detect viruses in animal species for which serological assays are not feasible due to the lack of species-specific secondary antibodies or for which cultured cells suitable for growth and titration of the virus are not available. For example, ranavirus in axolotls (*Ambystoma mexicanum*) and red-eared slider turtles (*Trachemys scripta elegans*)^{1,31} and monkey pox in giant pouched rats (*Cricetomys gambianus*) and black-tailed prairie dogs (*Cynomys ludovicianus*)⁶⁹ have all been detected using PCR.

Bacterial infections

Traditionally, bacterial infections have been diagnosed by culture in nutrient or selective media under aerobic, micro-aerophilic, or anaerobic conditions.^{24,63} Culturing of bacteria is best performed prior to the administration of antibiotics, and for slow-growing bacteria such as Mycobacteria, can take up to 6 wk. Collection of specimens from sites where normal flora is prevalent may result in overgrowth by normal, nonpathogenic bacteria and may mask the presence of pathogenic bacteria. If a sample is being screened for multiple bacteria, multiple types of culture media and culture conditions may be required. Once a bacterium has been cultured, bacterial smears can be stained to determine bacterial morphology. Panels of enzymatic and fermentation tests or Matrix-Assisted Laser Desorption/Ionization-Time Of Flight Mass Spectrometry (MALDI-TOF) can be performed on a bacterial isolate to speciate the bacteria. The antibiotic susceptibility of a bacterial isolate can also be determined. Bacterial infections can also be detected in stained tissue sections. For example, *Filobacterium rodentium* (previously: cilia-associated respiratory bacillus) and *Helicobacter* can be visualized in the respiratory or gastrointestinal tract using a silver based stain such as Warthin-Starry.^{56,134} Serologic methods are available for the detection of infections with some bacterial species.

Bacterial respiratory infections are difficult to detect using soiled bedding sentinels, as the bacteria usually are not present at sufficient levels in soiled bedding to cause infection and subsequent seroconversion of the sentinel. For example, *F. rodentium* is not effectively detected using soiled bedding sentinels, but can be detected by PCR in frozen or fixed respiratory tissues from infected rodents, or from oral or nasal swabs.^{29,30} PCR assays have been developed to detect a wide range of rodent bacterial agents, and PCR can be highly effective at differentiating pathogenic

from closely related nonpathogenic bacterial species. For example, biochemical testing of bacteria cultured from the skin of *C. bovis* infected nude mice may reveal several *Corynebacterium* species with the same biochemical profile, whereas MALDI-TOF or a PCR assay that targets the *rpoB* virulence gene of *C. bovis* will specifically detect *C. bovis*.^{37,59}

Not all bacteria can be cultured. Likewise, the different growth rates of the bacteria present in the sample can lead to overgrowth of faster-growing bacterial species, and thus obscure the detection of slower growing bacterial species. In addition, bacteria in the sample that require specialized media or growth conditions may not be detected if the specialized media or growth conditions are not used during screening. An excellent example of this limitation of culture as a diagnostic technique is *Helicobacter*. More than 10 *Helicobacter* species can infect rodents, and rodents can be infected with more than one *Helicobacter* species at the same time.¹³⁶ In addition, *Helicobacter* are fastidious bacteria that are difficult to culture. These factors create a challenge when determining the species of *Helicobacter* present in infected animals. In contrast, if PCR is used as the method of diagnosis, generic *Helicobacter* primers that detect most *Helicobacter* species can be used,¹²² or primers can be designed to detect only a single species, such as *Helicobacter hepaticus*.⁴⁹ Alternatively, multiple sets of primers can be used to detect several species of *Helicobacter* in a single assay (multiplex PCR).^{47,110} Multiplex PCR, like other multiplexed assays such as multiplexed fluorometric immunoassays, uses fewer reagents and less sample material than multiple single assays, and is therefore more cost-effective. When generic *Helicobacter* primers are used, the PCR product may then be sequenced to determine the *Helicobacter* species that was amplified. PCR also can be used to rapidly detect other bacteria that grow slowly in bacterial culture media, such as *Mycobacteria* and *Mycoplasma*.^{91,98,125} Even for bacteria that can be readily cultured, such as *Rodentibacter*, PCR is more sensitive and more rapid than culturing.^{13,112} For zoonotic bacteria, such as *Streptobacillus moniliformis*, using PCR rather than culture-based methods of detection has the additional benefit of decreasing the risk to laboratory personnel.⁵¹

Parasitic infections

Infestation of rodents with fur mites (*Myobia musculi*, *Myocoptes musculinus* and *Radfordia affinis*) has traditionally been diagnosed by visualizing adult mites, their larvae and nymphs, or their eggs using microscopy.³ Because transmission of fur mites via soiled bedding is unreliable, direct sampling of colony mice is recommended.⁸⁴ PCR of DNA extracted from fur swabs was more sensitive than fur pluck, tape, and sticky paper tests, and its sensitivity was similar to postmortem pelt exams.^{77,121,135} Fur mite PCR can also detect mite DNA at all stages of their life cycle. Because the rRNA genes of *M. musculis* and *R. affinis* are 99% homologous, both mites can be amplified using the same primers.⁶² In contrast, primers for *M. musculinus* amplify only *M. musculinus*.⁸³ PCR of feces has also been used to detect the presence of *M. musculinus* ingested by the mice.⁸³ The ability to detect *M. musculinus* infestation in a mouse colony using PCR was affected by the age of the mice tested, as older mice had lower egg and mite loads.¹²¹ For optimal detection, the sampling site on the mouse should be considered, as adult *M. musculinus* loads were highest in fur samples from the ventral abdomen and dorsal tail base but egg numbers were highest along the dorsal midline just caudal to the ears.⁹⁹ Because PCR detects DNA from both live and dead mites, positive fur swab PCR results from mice treated with antiparasitics can indicate a residual active infection as well as the presence of residual dead mites

on the mouse.¹²¹ Follicular mites such as *Demodex musculi* are generally considered to be commensal but can cause disease in immunocompromised mice.^{107,130} *D. musculi* has been detected in skin samples from immunocompromised mice using PCR.¹⁰⁸ Antemortem detection of *D. musculi* in immunocompetent mice is challenging, due to the very low mite load present in these animals. Both PCR-based and conventional methods of mite testing can fail to detect *D. musculi* in quarantined mice, leading to the introduction of *D. musculi* into established mouse colonies.¹⁰⁷

Rodent pinworms (*Aspicularis tetraptera*, *Syphacia obvelata* and *Syphacia muris*) traditionally have been diagnosed by examination of feces, intestinal contents or perianal tape samples using microscopy; these species can be distinguished morphologically.^{3,40} Pinworm PCR is highly specific for differentiating the 3 pinworm species, even when rodents are infected with more than one pinworm species.^{26,38,58,60,114} PCR of fecal DNA can also be used to detect *A. tetraptera* eggs and worms. Several studies have compared the sensitivity of fecal *A. tetraptera* PCR to fecal flotation, fecal centrifugation concentration or direct worm detection in intestinal contents, with divergent results.^{38,58,60} In some cases, fecal flotation, fecal centrifugation concentration or direct worm detection tests were positive for *A. tetraptera*, while *A. tetraptera* fecal PCR was negative. While hundreds of adult *A. tetraptera* can be present in the intestine of a single infested mouse, and each adult female worm can shed eggs several times during their 21 to 26 d long reproductive lifespan, the number of eggs present in the intestine, and therefore in the feces, can vary widely. Because the number of eggs isolated from feces was highest in 8 to 12 wk old mice, testing of feces from young adult mice is recommended when testing a colony for *A. tetraptera*.⁶⁰ PCR of fecal DNA can be used to detect *Syphacia obvelata* or *Syphacia muris* eggs and adult worms.^{26,58,114} Fecal *S. obvelata* PCR is more sensitive than either perianal tape tests or direct worm detection from the intestine.⁵⁸ Because adult female *S. obvelata* worms migrate to the anus and release their eggs in a single burst, false-negative tape tests can occur if testing occurs between periods of egg release.¹²⁷ The intermittent release of eggs can also lead to false-negative pinworm PCR results if feces are collected at times when eggs are not being shed. Repeated collection and testing of fecal samples may be necessary to confirm the presence or absence of pinworm infection. *Syphacia muris*, shows a 2 to 3 wk periodicity of egg shedding, with eggs shed in the highest numbers in midafternoon.⁹⁷

Other intestinal parasites such as *Entamoeba muris*, *Giardia muris*, *Spironucleus muris* and *Tritrichomonas muris* can be visualized in intestinal contents. PCR assays for these intestinal parasites have been developed and are routinely performed as part of quarantine procedures and vendor health screens.^{6,43,65,73,116}

Fungal infections

In general, fungi grow slowly in culture, and fungal infections often are diagnosed histologically. The most common fungal pathogens present in rodents, *Pneumocystis murina* and *Pneumocystis carinii*, have traditionally been diagnosed histologically from lung samples.³ PCR of lungs, bronchoalveolar lavages, nasopharyngeal aspirates and oral swabs can be used to detect *Pneumocystis spp.* DNA.^{46,70,88,113} *Batrachochytrium dendrobatidis* is a fungal pathogen of high concern in amphibians (anurans, urodeles, and caecilians). Skin swabs from axolotls (*Ambystoma mexicanum*) and rough-skinned newts (*Taricha granulosa*) have been used to detect *B. dendrobatidis* using PCR.³⁴ Pathogen-specific primers can be used when a particular fungal species is suspected. If a fungal infection is suspected in an animal and

no particular fungal species cultured or identified morphologically, PCR using generic fungal primers and/or genus-specific primers, followed by sequencing of the PCR product, can be used to speciate fungi that are detected.¹³⁷

Biologic contaminant testing

All samples (tumors, cell lines, etc.) to be inoculated into rodents should be tested for the presence of rodent viruses and *M. pulmonis*. Human patient derived xenograft tumors, tissues and stocks intended to be introduced into nude mice should also be tested for *C. bovis*.⁹⁵

The mouse antibody production (MAP) test was developed over 70 y ago and involves the inoculation of the biologic material into mice via several routes (intraperitoneally, intranasally and orally), followed several weeks later by the collection of blood from the mice for serological testing to detect the presence of infectious agents to be excluded.¹²³ MAP testing requires an area for isolation of the inoculated mice from the rest of the colony.

Once cell-culture based methods were developed for growing most of the agents to be excluded, cell-culture based testing also was used to detect biologic contaminants.^{7,12,33,89} This method involves inoculation of several susceptible cell lines with the test substance and then staining of the inoculated cells with antiviral antibodies to determine if the cells have produced viral antigens.

MAP testing and cell-culture methods have now been replaced by PCR/RT-PCR to detect viral and bacterial nucleic acids present in the cultured cells, tissue or /tumor samples. Molecular testing of biologic materials has equivalent or greater sensitivity than traditional MAP testing,^{12,14} and PCR-based biologic contaminant testing is faster than mouse or cell-culture based testing. PCR-based testing is also consistent with the principle of the 3Rs in that animals are replaced by an alternative method. PCR can be used to detect agents that cannot be readily grown in cell culture, such as lactate dehydrogenase virus.¹⁹ Furthermore, PCR testing does not result in the production of viruses in cells or animals that can infect rodents or humans; this is particularly important for lethal viruses such as lymphocytic choriomeningitis virus.⁹

PCR of samples collected noninvasively from rodents

Detecting infections using samples collected with noninvasive methods has several advantages. First, it likely provides better animal welfare than does invasive collection. Second, noninvasive collection methods can be performed more frequently than invasive methods without generating animal welfare concerns. Traditionally, performing multiple serological tests required more than 100 μ L of blood, which was collected from the retro-orbital sinus of anesthetized mice. This method can be performed weekly if the eyes are alternated. Newer commercial serology methods require less blood, such that unanesthetized mice can be bled from the facial, saphenous or tail vein; however, restraint is needed, and sedation, vasodilation, or shaving may be necessary.^{67,120}

In contrast, collection of feces, skin swabs, or fur swabs can be performed daily with gentle manual restraint of the rodent. The ability to collect multiple samples over a short period from the same mouse allows for the time course of agent shedding to be determined.^{22,68,78} Sequential samples can be used to determine when a mouse has cleared the infection and therefore is no longer a transmission risk to other colony animals.^{21,87} For

agents that are shed for only a short period of time (like murine rotavirus) or that are intermittently shed (such as pinworms), infection can be missed if only a single sample is collected. For this type of infectious agent, PCR should be used in conjunction with traditional methods (serology, bacteriology or parasitology). PCR of samples taken from individual colony mice for health monitoring is less cost effective because of the large number of samples that would have to be tested, as compared with serology or PCR based testing of sentinel mice. However, due to the high sensitivity of PCR, up to 10 fecal pellets or swabs can be pooled prior to nucleic acid extraction and still yield a positive result.⁹³ Therefore, all mice in a cage can be tested as a single sample. The effect of pooling samples on the sensitivity of PCR has not been tested for all agents, and pooling samples could complicate detection of agents that are shed poorly or intermittently. A single fecal sample can yield sufficient nucleic acids to test for a panel of more than 20 rodent intestinal viruses, bacteria, and parasites. Samples can also be pooled for other diagnostic methods (serology, culture, and microscopy).

The use of bacterial 16S rRNA gene primers can result in the amplification of several closely related bacterial species (such as multiple *Helicobacter* species) in fecal samples.¹²² In contrast, for detection of a single pathogenic species of bacteria in feces, primers specific for a virulence factor gene are often used. For example, primers specific for the *espB* gene of *Citrobacter rodentium* can be used for PCR testing of mouse feces.^{96,124}

Respiratory agents, such as Sendai virus, *F. rodentium* and *Rodentibacter pneumotropicus/Rodentibacter heylii* (previously *Pasteurella pneumotropica* biotypes Jawetz and Heyl⁸) have been difficult to detect using soiled bedding sentinels.^{2,32,101,102} Most likely, the amount of these agents that are shed into the bedding and remain infectious is very low. PCR of fecal samples can be used to detect respiratory agents because animals swallow respiratory secretions that contain infectious and noninfectious forms of the agent. However, the sensitivity of detection of respiratory agents in feces is less than that for intestinal agents.⁶⁵ Alternatively, minimally invasive oral or nasal swabs can be used to test for respiratory pathogens.^{25,52,61,65} A combination of feces, fur swabs, and oral swabs can be used to test quarantined mice for a panel of more than 30 rodent viruses, bacteria, fungi, and parasites.⁶⁵ The direct testing of quarantined mice using PCR is therefore consistent with the principle of the 3Rs, in that it eliminates the need for contact sentinels in quarantine and eliminates the production of unwanted litters when imported breeding pairs are housed with female sentinel mice.

PCR of environmental samples

The collection of environmental samples is even less stressful than the collection of feces or fur samples, as it does not require rodent handling. Culturing of infectious agents from environmental surfaces is of limited use, as desiccation drastically decreases the infectivity for many bacteria and viruses.⁵⁴ However, samples from environmental sites, including soiled bedding, cage filter tops and individually ventilated cage (IVC) racks, can be collected for PCR testing. One drawback of health monitoring using fecal or fur swab PCR is that it only detects infectious agents that are currently in or on the animal. The use of PCR of environmental samples as part of a health monitoring program allows the detection of agents deposited over a longer period, as each sample represents the accumulation of infectious agents and noninfectious debris deposited into the environment since the last time the sample site was cleaned. Consideration of what sample location to test, how frequently to test, and how

to disinfect the environment if an agent is detected is important when designing a PCR-based environmental monitoring program for infectious agents in research colonies.

Infectious agents shed via multiple routes (fecal, oral, respiratory, skin, fur) can be detected by PCR on cage surfaces and/or in soiled bedding. During an outbreak of rat coronavirus (shed in tears and by sneezing), cage surface swab RT-PCR was more sensitive than RT-PCR of Harderian and salivary glands and could be performed earlier during the infection than serology.²⁵ During an outbreak of mouse parvovirus (MPV), which is shed in feces, PCR of DNA extracted from swabs of the portion of the cage including the soiled bedding were more effective at detecting MPV in late infection than soiled bedding sentinels, but less effective than pooled fecal PCR.⁸⁶ Cage/soiled bedding swabs have also been used to detect fur mites.⁷⁴ In cages that contain mice infested/infested with several infectious agents (such as wild mice or those obtained from pet stores), cage swabs represent a single sample that can be tested for multiple agents.²⁶ Soiled bedding and nesting material can also be tested using PCR.^{16,138}

Several methods to monitor the dust present in air exhausted from IVC cages and racks have been developed. These methods can be used to monitor laboratory animal colonies for infectious agents, although differences in the airflow in different models of cages and racks may affect the sensitivity of the method. For IVC rack systems in which the air is filtered as it is exhausted from the cage, testing is best performed at the cage level. Monitoring at the cage level can be accomplished by placing pieces of gauze or filter paper on the underside of the lid of cages of colony or sentinel mice. PCR of nucleic acids extracted from the collection material after 1 to 90 d could detect MHV, MPV, murine norovirus (MNV), Sendai virus, *Helicobacter spp.*, *Rodentibacter spp.*, *M. muscoli*, *M. musculus*, *A. tetraptera*, *R. affinis*, *S. obvelata*, *Spiroplasma muris* and *E. muris*.^{23,29,57} Filters placed on sentinel cage lids for 3 mo, compared with 1 mo, had a higher rate of detection likely because they were exposed to 6 doses of soiled bedding rather than only 2 doses.³⁹ PCR of DNA extracted from filter top swabs was also effective at detecting fur mites, pinworms and *C. bovis*.^{57,95} A recent study showed that filter material placed on the underside of an IVC cage that did not contain sentinel mice but received soiled bedding every other week for 1 to 3 mo was effective at detecting MNV, *Helicobacter spp.*, *Rodentibacter spp.*, *Spiroplasma muris* and *E. muris* if the cages were shaken for 15 seconds twice weekly to aerosolize small particles from the soiled bedding.³⁹

For IVC rack systems in which the exhaust air is filtered at the rack level, exhaust air dust can be monitored by PCR of filter material placed on the exhaust prefilter, swabs of the exhaust plenum, or filters placed in the exhaust air stream. Several studies have investigated IVC exhaust prefilter testing using several types of IVC racks, durations of filter placement and prevalence of the infectious agents on the IVC rack.^{6,23,79,102,138} PCR of nucleic acids extracted from gauze placed on the IVC exhaust prefilter for 1 to 42 d was effective at detecting MHV, MPV, Sendai virus, and *Helicobacter spp.* shed from 4 cages of mice experimentally infected with each agent.²³ PCR of nucleic acids extracted from gauze placed on the IVC exhaust prefilter for 1 to 12 wk detected MHV, MPV, *Helicobacter spp.*, *Rodentibacter spp.*, *E. muris*, *T. muris*, *M. musculus*, *R. affinis*, *A. tetraptera* and *S. obvelata*, but not MNV, shed from 1 to 5 cages of mice infected with each agent.⁶ PCR of nucleic acids extracted from pieces of gauze placed on the IVC exhaust prefilter for 6 or 12 wk were able to detect several endemic murine viruses and bacteria (MNV, *Rodentibacter spp.*, *Helicobacter sp.*, *Klebsiella oxytoca*, and

Proteus mirabilis) even when the prevalence of these agents was low in the mice on the rack.^{102,138} PCR of nucleic acids extracted from pieces of gauze placed on the IVC exhaust prefilter for 1 or 2 wk were able to detect endemic infection with MNV and *M. musculus*.⁷⁹

Several studies have investigated using exhaust manifold swabs to monitor exhaust air dust from a variety of IVC racks in which the exhaust air is not filtered at the cage level. Fur mite (*M. muscoli* or *R. affinis*) DNA were detected on the horizontal exhaust manifolds of most IVC racks within 2 wk if the rack housed even a single cage of fur mite infested mice.⁷⁴ *C. bovis* DNA was detected on the horizontal exhaust manifolds of IVC racks within 1 wk when each rack housed only one cage of *C. bovis* infected mice.⁹⁴ *A. tetraptera* DNA was detected in swabs taken from 4 locations in the air handling unit and IVC rack one week after housing multiple cages of pinworm-infested mice.⁷⁶ In a recent study, over 1700 IVC exhaust plenum swabs were tested by PCR for 14 opportunistic bacteria and *P. murina*; *Staphylococcus xylosum*, *Staphylococcus aureus*, *R. pneumotropicus*, *R. heylia*, *Proteus mirabilis*, *Pseudomonas aeruginosa*, *Klebsiella oxytoca* and *Klebsiella pneumoniae* were detected in at least 1% of the swabs.¹¹⁸

The duration of exhaust air dust exposure needed to detect an agent seems to depend on the amount of infectious agent released into the airflow. A shorter time is necessary if most of the cages on the rack house infected/infested mice or during acute infections when high levels of the agent are being produced. Several IVC rack manufacturers have designed their racks to facilitate exhaust air dust monitoring through the incorporation of specially designed filters and filter holders. A recent study compared 2 of these systems and showed that *Entamoeba spp.*, *Helicobacter spp.* and *Rodentibacter spp.* could be detected with either filter system.⁹⁰ In another study, exhaust air dust monitoring using the specially designed filters was more sensitive than soiled bedding sentinels at detecting endemic infections with *Helicobacter spp.* and *Rodentibacter spp.*⁹² An essential aspect of using PCR testing of IVC components is complete decontamination of the rack prior to additional testing if an agent is detected, as carryover of nucleic acids can lead to false positive results. False positives can also occur if the primers used will amplify closely related species, such as occurred with the detection of rhabditid nematodes in unautoclaved corn cob bedding using *Aspiculuris* primers.⁸²

Any location in the animal facility in which debris generated by laboratory animals accumulates can be tested for infectious agent nucleic acids. PCR of debris collected from the prefilter of an animal bedding disposal cabinet detected several infectious agents endemic in the mouse colony (MNV and *Helicobacter spp.*) and several infectious agents present in a small number of cages of experimentally infected mice (MPV, *M. pulmonis*, *S. obvelata*, and *M. musculus*).²⁶ After vaporized hydrogen peroxide decontamination of a facility that had housed mice infected with *C. bovis*, PCR of swabs of equipment and surfaces in the room and in the adjacent corridors were used to validate the decontamination process.¹⁰⁰ For aquatic species, aquarium water or sediment can be tested by PCR to monitor for DNA of infectious agents. PCR of sump swabs were more sensitive than PCR of zebrafish for detecting mycobacteria.¹⁰⁴ *Pseudoloma neurophilia* DNA was detected in eggs and sperm from zebrafish and from aquarium water if the sample was sonicated to disrupt the spores.¹²⁶ PCR testing of filters for water and sediment from aquariums housing zebrafish detected *Mycobacterium spp.*, *Myxidium streisingeri*, *P. neurophilia* and *Pseudocapillaria tomentosa*, with *Mycobacterium spp.* detected most frequently.^{28,103} *Pseudocapillaria xenopi* was

detected by PCR of sediment from aquariums housing African clawed frogs.⁴⁵

Germ-free Animals and Microbiome Analysis

In addition to routine health monitoring and diagnostic testing of immunocompetent and immunodeficient mice that are colonized with a wide range of commensal organisms, many laboratory animal facilities also maintain colonies of gnotobiotic (including germ-free) animals that require specialized health monitoring. PCR using bacterial 16S rRNA primers, in addition to bacterial culture, are used to confirm the absence of bacteria in germ-free animals.⁵⁰ PCR on feces from germ-free mice can reveal traces of bacterial DNA from natural-ingredient based diets, even after the diet is autoclaved, giving the impression that the animal and/or the isolator have been compromised. Therefore, PCR alone should not be used to determine if a biosecurity breach has occurred.

One use of germ-free animals is to study the role of different microbiomes in animal models of disease such as cancer, colitis and diabetes.⁶⁴ The microbiome of germ-free animals can be reconstituted with a single bacterial agent or with a complex mix of bacterial agents to study how these agents affect the disease phenotype. Both husbandry (housing, diet, bedding) and animal (genotype, age, sex and source) factors can alter the mouse fecal microbiome.^{11,17,20,35,41,42,105,111,115,119,131,132} While microbiome analysis is currently not routinely used as a diagnostic tool in laboratory animal medicine, an understanding of how microbiomes are analyzed and what can alter the microbiome is essential to the field.²⁷ One method for analyzing the microbiome uses 16S rRNA PCR, followed by sequencing of the pool of PCR products generated.⁶⁶ The newer shotgun metagenomics sequencing method does not require PCR, and instead involves breaking the microbial DNA present in the sample into small fragments for sequencing.⁷⁵ Bioinformatics analysis of the millions of sequences generated by these methods allows determination of the relative abundance of each microbial species in the population. As sequencing of the complete microbiome (bacteria, viruses, fungi, and parasites) become more affordable, this methodology may replace current diagnostic methods for identifying known infectious agents in rodents and other laboratory animals and has the potential to identify novel disease-causing agents.

Conclusions

PCR has many uses in laboratory animal science and medicine. In some situations, PCR-based testing has replaced other methods, whereas in other cases, using PCR in conjunction with other methods improves the ability to monitor for infectious agents in research animal colonies and their environment. The high specificity of PCR is advantageous in that it can be used to differentiate closely related nonpathogenic and pathogenic strains of an infectious agent. However, small changes in the primer sites of new strains of an agent can lead to false negatives. The second advantage of PCR is its sensitivity (the ability to detect only a few infectious agents); however, this high sensitivity can produce false positives if extraction or PCR reagents become contaminated with low levels of extraneous nucleic acids. Molecular diagnostics are rapid and, when used in the face of an outbreak, can quickly detect which animals are shedding the agent and pose a risk for transmission to other animals in the facility. PCR can be used on a wide range of species, tissue types, excreta and environment samples. An important facet of using PCR is knowing where in or on the animal the infectious agent is most likely to be found and collecting samples

from the animal only during active infection. Conversely, PCR testing of environmental samples reveals both current and past infections, as it detects all agents deposited at the sampling site since the site was last sanitized. PCR detects a wide range of infectious agents (viruses, bacteria, fungi, and parasites) and can be used to detect agents that cannot be cultured or that are poorly transmitted to sentinels, such as respiratory agents. The minimal invasiveness of sampling and the reduction in number of animals needed and sampling procedures are all consistent with the ethical principles of the 3Rs.

Acknowledgment

This manuscript is based on the Wallace P Rowe lecture presented at the 2019 National Meeting of the American Association for Laboratory Science.

References

1. Allender MC, Mitchell MA, Tores T, Sekowska J, Driskell EA. 2013. Pathogenicity of frog virus 3-like virus in red-eared turtles (*Trachemys scripta elegans*) at two environmental temperatures. *J Comp Pathol* 149:356–367. <https://doi.org/10.1016/j.jcpa.2013.01.007>.
2. Artwohl JE, Cera LM, Wright MF, Medina LV, Kim LJ. 1994. The efficacy of a dirty bedding sentinel system for detecting Sendai virus infection in mice: A comparison of clinical signs and seroconversion. *Lab Anim Sci* 44:73–75.
3. Baker DG. 2003. Pathogens of rats and mice, p 19–107. In: Natural pathogens of laboratory animals: their effects on research. Washington (DC): ASM Press.
4. Balansard I, Cleverley L, Cutler KL, Spangberg MG, Thibault-Duprey K, Langermans JA. 2019. Revised recommendations for health monitoring of non-human primate colonies (2018): FELASA working group report. *Lab Anim* 53:429–446. <https://doi.org/10.1177/0023677219844541>.
5. Barthold SW, Smith AL, Povar ML. 1985. Enterotropic mouse hepatitis virus infection in nude mice. *Lab Anim Sci* 35:613–618.
6. Bauer BA, Besch-Williford CL, Livingston RS, Crim MJ, Riley LK, Myles MH. 2016. Influence of rack design and disease prevalence on detection of rodent pathogens in exhaust air debris samples from individually ventilated caging systems. *J Am Assoc Lab Anim Sci* 55:782–788.
7. Bauer BA, Besch-Williford CL, Riley LK. 2004. Comparison of the mouse antibody production (MAP) assay and polymerase chain reaction (PCR) assays for the detection of viral contaminants. *Biologicals* 32:177–182. <https://doi.org/10.1016/j.biologics.2004.08.004>.
8. Benga L, Sager M, Christensen H. 2018. From the [*Pasteurella*] *pneumotropica* complex to *Rodentibacter* spp.: An update on [*Pasteurella*] *pneumotropica*. *Vet Microbiol* 217:121–134. <https://doi.org/10.1016/j.vetmic.2018.03.011>.
9. Besselsen DG, Wagner AM, Loganbill JK. 2003. Detection of lymphocytic choriomeningitis virus by use of fluorogenic nuclease reverse transcriptase-polymerase chain reaction analysis. *Comp Med* 53:65–69.
10. Bhatt PN, Percy DH, Jonas AM. 1972. Characterization of the virus of sialodacryoadenitis of rats: A member of the coronavirus group. *J Infect Dis* 126:123–130. <https://doi.org/10.1093/infdis/126.2.123>.
11. Bidot WA, Ericsson AC, Franklin CL. 2018. Effects of water decontamination methods and bedding material on the gut microbiota. *PLoS One* 13:1–16. <https://doi.org/10.1371/journal.pone.0198305>.
12. Blank WA, Henderson KS, White LA. 2004. Virus PCR assay panels: An alternative to the mouse antibody production test. *Lab Anim (NY)* 33:26–32. <https://doi.org/10.1038/labam0204-26>.
13. Bootz F, Kirschnek S, Nicklas W, Wyss SK, Homberger FR. 1998. Detection of *Pasteurellaceae* in rodents by polymerase chain reaction analysis. *Lab Anim Sci* 48:542–546.
14. Bootz F, Sieber I, Popovic D, Tischhauser M, Homberger FR. 2003. Comparison of the sensitivity of in vivo antibody production tests with in vitro PCR-based methods to detect infectious

- contamination of biological materials. *Lab Anim* 37:341–351. <https://doi.org/10.1258/002367703103051895>.
15. **Bruggeman CA, Debie WM, Grauls G, Majoor G, van Boven CP.** 1983. Infection of laboratory rats with a new cytomegalo-like virus. *Arch Virol* 76:189–199. <https://doi.org/10.1007/BF01311103>.
 16. **Buchheister S, Roegener F, Zschemisch N, Talbot SR, Christensen H, Bleich A.** 2019. One for two: A novel and highly sensitive virulence factor-based quantitative polymerase chain reaction assay for the simultaneous detection of *Rodentibacter pneumotropicus* and *Rodentibacter heylii* in environmental sample material. *Lab Anim* 54:239–250. <https://doi.org/10.1177/0023677219853600>.
 17. **Campbell JH, Foster CM, Vishnivetskaya T, Campbell AG, Yang ZK, Wymore A, Palumbo AV, Chesler EJ, Podar M.** 2012. Host genetic and environmental effects on mouse intestinal microbiota. *ISME J* 6:2033–2044. <https://doi.org/10.1038/ismej.2012.54>.
 18. **Charles River.** 2020. PCR infectious agent testing. [Cited 13 January 2020]. Available at: <https://www.criver.com/products-services/research-models-services/animal-health-surveillance/pcr-infectious-agent-testing>.
 19. **Chen Z, Plagemann PG.** 1997. Detection of lactate dehydrogenase-elevating virus in transplantable mouse tumors by biological assay and RT-PCR assays and its removal from the tumor cell. *J Virol Methods* 65:227–236. [https://doi.org/10.1016/S0166-0934\(97\)02188-5](https://doi.org/10.1016/S0166-0934(97)02188-5).
 20. **Choo JM, Trim PJ, Leong LEX, Abell GCJ, Brune C, Jeffries N, Wesselingh S, Dear TN, Snel MF, Rogers GB.** 2017. Inbred mouse populations exhibit intergenerational changes in intestinal microbiota composition and function following introduction to a facility. *Front Microbiol* 8:1–14. <https://doi.org/10.3389/fmicb.2017.00608>.
 21. **Compton SR, Ball-Goodrich LJ, Paturzo FX, Macy JD.** 2004. Transmission of enterotropic mouse hepatitis virus from immunocompetent and immunodeficient mice. *Comp Med* 54:29–35.
 22. **Compton SR, Booth CJ, Macy JD.** 2017. Murine astrovirus infection and transmission in neonatal CD1 mice. *J Am Assoc Lab Anim Sci* 56:402–411.
 23. **Compton SR, Homberger FR, Paturzo FX, Clark JM.** 2004. Efficacy of three microbiological monitoring methods in a ventilated cage rack. *Comp Med* 54:382–392.
 24. **Compton SR, Riley LK.** 2001. Detection of infectious agents in laboratory rodents: Traditional and molecular techniques. *Comp Med* 51:113–119.
 25. **Compton SR, Vivas-Gonzalez BE, Macy JD.** 1999. Reverse transcriptase polymerase chain reaction-based diagnosis and molecular characterization of a new rat coronavirus strain. *Lab Anim Sci* 49:506–513.
 26. **Compton SR, Macy JD.** 2015. Effect of cage-wash temperature on the removal of infectious agents from caging and the detection of infectious agents on the filters of animal bedding-disposal cabinets by PCR analysis. *J Am Assoc Lab Anim Sci* 54:745–755.
 27. **Costa M, Weese JS.** 2019. Methods and basic concepts for microbiota assessment. *Vet J* 249:10–15. <https://doi.org/10.1016/j.tvjl.2019.05.005>.
 28. **Crim MJ, Lawrence C, Livingston RS, Rakitin A, Hurley SJ, Riley LK.** 2017. Comparison of antemortem and environmental samples for zebrafish health monitoring and quarantine. *J Am Assoc Lab Anim Sci* 56:412–424.
 29. **Cundiff DD, Besch-Williford C, Hook RRJ, Franklin CL, Riley LK.** 1994. Detection of cilia-associated respiratory bacillus by PCR. *J Clin Microbiol* 32:1930–1934. <https://doi.org/10.1128/JCM.32.8.1930-1934.1994>.
 30. **Cundiff DD, Riley LK, Franklin CL, Hook RRJ, Besch-Williford C.** 1995. Failure of a soiled bedding sentinel system to detect cilia-associated respiratory bacillus infection in rats. *Lab Anim Sci* 45:219–221.
 31. **Davidson EW, Jancovich JK, Borland S, Newberry M, Gresens J.** 2003. What's your diagnosis?: Dermal lesions, hemorrhage, and limb swelling in laboratory axolotls. *Lab Anim* 32:23–25. <https://doi.org/10.1038/lablan0303-23>.
 32. **de Bruin WCC, van de Ven EME, Hooijmans CR.** 2016. Efficacy of soiled bedding transfer for transmission of mouse and rat infections to sentinels: A systematic review. *PLoS One* 11:1–11. <https://doi.org/10.1371/journal.pone.0158410>.
 33. **de Souza M, Smith AL.** 1989. Comparison of isolation in cell culture with conventional and modified mouse antibody production tests for detection of murine viruses. *J Clin Microbiol* 27:185–187. <https://doi.org/10.1128/JCM.27.1.185-187.1989>.
 34. **Del Valle JM, Eisthen HL.** 2019. Treatment of chytridiomycosis in laboratory axolotls (*Ambystoma mexicanum*) and rough-skinned newts (*Taricha granulosa*). *Comp Med* 69:204–211. <https://doi.org/10.30802/AALAS-CM-18-000090>.
 35. **Dias MF, Reis MP, Acurcio LB, Carmo AO, Diamantino CF, Motta AM, Kalapothakis E, Nicoli JR, Nascimento AMA.** 2018. Changes in mouse gut bacterial community in response to different types of drinking water. *Water Res* 132:79–89. <https://doi.org/10.1016/j.watres.2017.12.052>.
 36. **Dillehay DL, Lehner ND, Huerkamp MJ.** 1990. The effectiveness of a microisolator cage system and sentinel mice for controlling and detecting MHV and Sendai virus infections. *Lab Anim Sci* 40:367–370.
 37. **Dole VS, Henderson KS, Fister RD, Pietrowski MT, Maldonado G, Clifford CB.** 2013. Pathogenicity and genetic variation of 3 strains of *Corynebacterium bovis* in immunodeficient mice. *J Am Assoc Lab Anim Sci* 52:458–466.
 38. **Dole VS, Zaias J, Kyricopoulos-Cleasby DM, Banu LA, Waterman LL, Sanders K, Henderson KS.** 2011. Comparison of traditional and PCR methods during screening for and confirmation of *Aspicularis tetraptera* in a mouse facility. *J Am Assoc Lab Anim Sci* 50:904–909.
 39. **Dubelko AR, Zuwannin M, McIntee SC, Livingston RS, Foley PL.** 2018. PCR testing of filter material from IVC lids for microbial monitoring of mouse colonies. *J Am Assoc Lab Anim Sci* 57:477–482. <https://doi.org/10.30802/AALAS-JAALAS-18-000008>.
 40. **Effler JC, Hickman-Davis JM, Erwin JG, Cartner SC, Schoeb TR.** 2008. Comparison of methods for detection of pinworms in mice and rats. *Lab Anim (NY)* 37:210–215. <https://doi.org/10.1038/lablan0508-210>.
 41. **Ericsson AC, Davis JW, Spollen W, Bivens N, Givan S, Hagan CE, McIntosh M, Franklin CL.** 2015. Effects of vendor and genetic background on the composition of the fecal microbiota of inbred mice. *PLoS One* 10:1–19. <https://doi.org/10.1371/journal.pone.0116704>.
 42. **Ericsson AC, Gagliardi J, Bouhan D, Spollen WG, Givan SA, Franklin CL.** 2018. The influence of caging, bedding, and diet on the composition of the microbiota in different regions of the mouse gut. *Sci Rep* 8:1–13. <https://doi.org/10.1038/s41598-018-21986-7>.
 43. **Fain MA, Karjala Z, Perdue KA, Copeland MK, Cheng LI, Elkins WR.** 2008. Detection of *Spiroplasma muris* in unpreserved mouse tissue and fecal samples by using a PCR assay. *J Am Assoc Lab Anim Sci* 47:39–43.
 44. **FELASA working group on revision of guidelines for health monitoring of rodents and rabbits, Mahler Convenor M, Berard M, Feinstein R, Gallagher A, Illgen-Wilcke B, Pritchett-Corning K, Raspa M.** 2014. FELASA recommendations for the health monitoring of mouse, rat, hamster, guinea pig and rabbit colonies in breeding and experimental units. *Lab Anim* 48:178–192.
 45. **Feldman SH, Ramirez MP.** 2014. Molecular phylogeny of *Pseudocapillaroides xenopi* (Moravec et Cosgrove 1982) and development of a quantitative PCR assay for its detection in aquarium sediment. *J Am Assoc Lab Anim Sci* 53:668–674.
 46. **Feldman SH, Weisbroth SP, Weisbroth SH.** 1996. Detection of pneumocystis carinii in rats by polymerase chain reaction: Comparison of lung tissue and bronchoalveolar lavage specimens. *Lab Anim Sci* 46:628–634.
 47. **Feng S, Ku K, Hodzic E, Lorenzana E, Freet K, Barthold SW.** 2005. Differential detection of five mouse-infecting *Helicobacter* species by multiplex PCR. *Clin Diagn Lab Immunol* 12:531–536. <https://doi.org/10.1128/CDLI.12.4.531-536.2005>.
 48. **Fenner F.** 1949. Mouse-pox; infectious ectromelia of mice; a review. *J Immunol* 63:341–373.
 49. **Fernér C, Lindberg AV, Feinstein RE.** 2002. Development and use of a simple polymerase chain reaction assay to screen for *Helicobacter* spp. and *H. hepaticus* in intestinal and fecal samples from laboratory mice. *Comp Med* 52:518–522.

50. Fontaine CA, Skorupski AM, Vowles CJ, Anderson NE, Poe SA, Eaton KA. 2015. How free of germs is germ-free? Detection of bacterial contamination in a germ-free mouse unit. *Gut Microbes* 6:225–233. <https://doi.org/10.1080/19490976.2015.1054596>.
51. Fornefett J, Krause J, Klose K, Fingas F, Hassert R, Eisenberg T, Schrödl W, Grunwald T, Müller U, Baums CG. 2018. Comparative analysis of clinics, pathologies and immune responses in BALB/c and C57BL/6 mice infected with *Streptobacillus moniliformis*. *Microbes Infect* 20:101–110. <https://doi.org/10.1016/j.micinf.2017.10.001>.
52. Franklin CL, Pletz JD, Riley LK, Livingston BA, Hook RRJ, Besch-Williford CL. 1999. Detection of cilia-associated respiratory (CAR) bacillus in nasal-swab specimens from infected rats by use of polymerase chain reaction. *Lab Anim Sci* 49:114–117.
53. Fredricks DN, Relman DA. 1999. Application of polymerase chain reaction to the diagnosis of infectious diseases. *Clin Infect Dis* 29:475–488. <https://doi.org/10.1086/598618>.
54. Gaertner DJ, Compton SR, Winograd DF. 1993. Environmental stability of rat coronaviruses (RCVs). *Lab Anim Sci* 43:403–404.
55. Gaertner DJ, Otto G, Batchelder M. 2007. Health delivery and quality assurance programs for mice, p 385–407. In: Fox JG, Barthold SW, Davison MT, Newcomer CE, editors. *The mouse in biomedical research*. Burlington (MA): Academic Press
56. Ganaway JR, Spencer TH, Moore TD, Allen AM. 1985. Isolation, propagation, and characterization of a newly recognized pathogen, cilia-associated respiratory bacillus of rats, an etiological agent of chronic respiratory disease. *Infect Immun* 47:472–479. <https://doi.org/10.1128/IAI.47.2.472-479.1985>.
57. Gerwin PM, Ricart Arbona RJ, Riedel ER, Henderson KS, Lipman NS. 2017. PCR testing of IVC filter tops as a method for detecting murine pinworms and fur mites. *J Am Assoc Lab Anim Sci* 56:752–761.
58. Gerwin PM, Ricart Arbona RJ, Riedel ER, Lephherd ML, Henderson KS, Lipman NS. 2017. Evaluation of traditional and contemporary methods for detecting *Syphacia obvelata* and *Aspiculuris tetraptera* in laboratory mice. *J Am Assoc Lab Anim Sci* 56:32–41.
59. Gonçalves JL, Tomazi T, Barreiro JR, Braga PA, Ferreira CR, Araújo Junior JP, Eberlin MN, dos Santos MV. 2014. Identification of *Corynebacterium spp.* isolated from bovine intramammary infections by matrix-assisted laser desorption ionization-time of flight mass spectrometry. *Vet Microbiol* 173:147–151. <https://doi.org/10.1016/j.vetmic.2014.06.028>.
60. Goodroe AE, Baxter VK, Watson J. 2016. Guidance regarding sample collection and refinement of fecal flotation exam for the isolation of *Aspiculuris tetraptera*. *J Am Assoc Lab Anim Sci* 55:541–547.
61. Goto K, Nozu R, Takakura A, Matsushita S, Itoh T. 1995. Detection of cilia-associated respiratory bacillus in experimentally and naturally infected mice and rats by the polymerase chain reaction. *Exp Anim* 44:333–336. <https://doi.org/10.1538/expanim.44.333>.
62. Grove KA, Smith PC, Booth CJ, Compton SR. 2012. Age-associated variability in susceptibility of Swiss Webster mice to MPV and other excluded murine pathogens. *J Am Assoc Lab Anim Sci* 51:789–796.
63. Hansen AK, Nielsen DS. 2015. *Handbook of laboratory animal bacteriology*, 2nd ed. Boca Raton (FL): CRC Press.
64. Hansen AK, Hansen CHF, Krych L, Nielsen DS. 2014. Impact of the gut microbiota on rodent models of human disease. *World J Gastroenterol* 20:17727–17736. <https://doi.org/10.3748/wjg.v20.i47.17727>.
65. Henderson KS, Perkins CL, Havens RB, Kelly ME, Francis BC, Dole VS, Shek WR. 2013. Efficacy of direct detection of pathogens in naturally infected mice by using a high-density PCR array. *J Am Assoc Lab Anim Sci* 52:763–772.
66. Hiegeist A, Glasner J, Reischl U, Gessner A. 2015. Analyses of intestinal microbiota: Culture versus sequencing. *ILAR J* 56:228–240. <https://doi.org/10.1093/ilar/ilv017>.
67. Hoggatt J, Hoggatt AF, Tate TA, Fortman J, Pelus LM. 2016. Bleeding the laboratory mouse: Not all methods are equal. *Exp Hematol* 44:132–137. <https://doi.org/10.1016/j.exphem.2015.10.008>.
68. Hsu CC, Riley LK, Wills HM, Livingston RS. 2006. Persistent infection with and serologic cross-reactivity of three novel murine noroviruses. *Comp Med* 56:247–251.
69. Hutson CL, Lee KN, Abel J, Carroll DS, Montgomery JM, Olson VA, Li Y, Davidson W, Hughes C, Dillon M, Spurlock P, Kazmierczak JJ, Austin C, Miser L, Sorhage FE, Howell J, Davis JP, Reynolds MG, Braden Z, Karem KL, Damon IK, Regnery RL. 2007. Monkeypox zoonotic associations: Insights from laboratory evaluation of animals associated with the multi-state US outbreak. *Am J Trop Med Hyg* 76:757–768. <https://doi.org/10.4269/ajtmh.2007.76.757>.
70. Icenhour CR, Rebholz SL, Collins MS, Cushion MT. 2001. Widespread occurrence of *Pneumocystis carinii* in commercial rat colonies detected using targeted PCR and oral swabs. *J Clin Microbiol* 39:3437–3441. <https://doi.org/10.1128/JCM.39.10.3437-3441.2001>.
71. IDEXX Bioanalytics. 2020. Real-time PCR. [Cited 13 January 2020]. Available at: <https://www.idexxbioanalytics.com/real-time-pcr>.
72. Ike F, Bourgade F, Ohsawa K, Sato H, Morikawa S, Saijo M, Kurane I, Takimoto K, Yamada YK, Jaubert J, Berard M, Nakata H, Hiraiwa N, Mekada K, Takakura A, Itoh T, Obata Y, Yoshiki A, Montagutelli X. 2007. Lymphocytic choriomeningitis infection undetected by dirty-bedding sentinel monitoring and revealed after embryo transfer of an inbred strain derived from wild mice. *Comp Med* 57:272–281.
73. Jackson GA, Livingston RS, Riley LK, Livingston BA, Franklin CL. 2013. Development of a PCR assay for the detection of *Spiro-nucleus muris*. *J Am Assoc Lab Anim Sci* 52:165–170.
74. Jensen ES, Allen KP, Henderson KS, Szabo A, Thulin JD. 2013. PCR testing of a ventilated caging system to detect murine fur mites. *J Am Assoc Lab Anim Sci* 52:28–33.
75. Jovel J, Patterson J, Wang W, Hotte N, O’Keefe S, Mitchel T, Perry T, Kao D, Mason AL, Madsen KL, Wong GK. 2016. Characterization of the gut microbiome using 16S or shotgun metagenomics. *Front Microbiol* 7:1–17. <https://doi.org/10.3389/fmicb.2016.00459>.
76. Kapoor P, Hayes YO, Jarrell LT, Bellinger DA, Thomas RD, Lawson GW, Arkema JD, Fletcher CA, Nielsen JN. 2017. Evaluation of anthelmintic resistance and exhaust air dust PCR as a diagnostic tool in mice enzootically infected with *Aspiculuris tetraptera*. *J Am Assoc Lab Anim Sci* 56:273–289.
77. Karlsson EM, Pearson LM, Kuzma KM, Burkholder TH. 2014. Combined evaluation of commonly used techniques, including PCR, for diagnosis of mouse fur mites. *J Am Assoc Lab Anim Sci* 53:69–73.
78. Kelmenson JA, Pomerleau DP, Griffey S, Zhang W, Karolak MJ, Fahey JR. 2009. Kinetics of transmission, infectivity, and genome stability of two novel mouse norovirus isolates in breeding mice. *Comp Med* 59:27–36.
79. Körner C, Miller M, Brielmeier M. 2019. Detection of murine astrovirus and *Myocoptes musculinus* in individually ventilated caging systems: Investigations to expose suitable detection methods for routine hygienic monitoring. *PLoS One* 14:1–12. <https://doi.org/10.1371/journal.pone.0221118>.
80. Kwok S, Higuchi R. 1989. Avoiding false positives with PCR. *Nature* 339:237–238. <https://doi.org/10.1038/339237a0>.
81. La Regina M, Woods L, Klender P, Gaertner DJ, Paturzo FX. 1992. Transmission of sialodacryoadenitis virus (SDAV) from infected rats to rats and mice through handling, close contact, and soiled bedding. *Lab Anim Sci* 42:344–346.
82. Leblanc M, Berry K, Graciano S, Becker B, Reuter JD. 2014. False-positive results after environmental pinworm PCR testing due to rhabditid nematodes in corncob bedding. *J Am Assoc Lab Anim Sci* 53:717–724.
83. Lee MA, Shen Z, Holcombe HR, Ge Z, Franklin EG, Ricart Arbona RJ, Lipman NS, Fox JG, Sheh A. 2019. Detection of *Myocoptes musculinus* in fur swab and fecal samples by using PCR analysis. *J Am Assoc Lab Anim Sci* 58:796–801. <https://doi.org/10.30802/AALAS-JAALAS-19-000046>.
84. Lindstrom KE, Carbone LG, Kellar DE, Mayorga MS, Wilkerson JD. 2011. Soiled bedding sentinels for the detection of fur mites in mice. *J Am Assoc Lab Anim Sci* 50:54–60.

85. Macy JD, Cameron GA, Smith PC, Ferguson TA, Compton SR. 2011. Detection and control of mouse parvovirus. *J Am Assoc Lab Anim Sci* 50:516–522.
86. Macy JD, Paturzo FX, Ball-Goodrich LJ, Compton SR. 2009. A PCR-based strategy for detection of mouse parvovirus. *J Am Assoc Lab Anim Sci* 48:263–267.
87. Macy JD, Smith PC, Wilson J, Compton SR. 2008. Mouse parovirus infection in immunodeficient mice. PS10. Abstract presented at the 59th AALAS National Meeting, Indianapolis, Indiana, 9–13 November 2008. *J Am Assoc Lab Anim Sci* 47:75.
88. Macy JD Jr, Weir EC, Compton SR, Shlomchik MJ, Brownstein DG. 2000. Dual infection with *Pneumocystis carinii* and *Pasteurella pneumotropica* in B cell-deficient mice: Diagnosis and therapy. *Comp Med* 50:49–55.
89. Mahabir E, Jacobsen K, Brielmeier M, Peters D, Needham J, Schmidt J. 2004. Mouse antibody production test: Can we do without it? *J Virol Methods* 120:239–245. <https://doi.org/10.1016/j.jviromet.2004.05.006>.
90. Mahabir E, Durand S, Henderson KS, Hardy P. 2018. Comparison of two prevalent individually ventilated caging systems for detection of murine infectious agents via exhaust air particles. *Lab Anim* 53:84–88. <https://doi.org/10.1177/0023677218785929>.
91. Mähler M, Bedigian HG, Burgett BL, Bates RJ, Hogan ME, Sundberg JP. 1998. Comparison of four diagnostic methods for detection of *Helicobacter* species in laboratory mice. *Lab Anim Sci* 48:85–91.
92. Mailhoit D, Ostdiek AM, Luchins KR, Bowers CJ, Theriault B, Langan GP. 2020. Comparing mouse health monitoring between soiled-bedding sentinel and exhaust air dust surveillance programs. *J Am Assoc Lab Anim Sci* 59:58–66. <https://doi.org/10.30802/AALAS-JAALAS-19-000061>.
93. Manuel CA, Hsu CC, Riley LK, Livingston RS. 2008. Soiled-bedding sentinel detection of murine norovirus 4. *J Am Assoc Lab Anim Sci* 47:31–36.
94. Manuel CA, Pugazhenthil U, Leszczynski JK. 2016. Surveillance of a ventilated rack system for *Corynebacterium bovis* by sampling exhaust-air manifolds. *J Am Assoc Lab Anim Sci* 55:58–65.
95. Manuel CA, Pugazhenthil U, Spiegel SP, Leszczynski JK. 2017. Detection and elimination of *Corynebacterium bovis* from barrier rooms by using an environmental sampling surveillance program. *J Am Assoc Lab Anim Sci* 56:202–209.
96. McKeel R, Douris N, Foley PL, Feldman SH. 2002. Comparison of an espB gene fecal polymerase chain reaction assay with bacteriologic isolation for detection of *Citrobacter rodentium* infection in mice. *Comp Med* 52:439–444.
97. Meade TM, Watson J. 2014. Characterization of rat pinworm (*Syphacia muris*) epidemiology as a means to increase detection and elimination. *J Am Assoc Lab Anim Sci* 53:661–667.
98. Meritet DM, Mulrooney DM, Kent ML, Löhr CV. 2017. Development of quantitative real-time PCR assays for postmortem detection of *Mycobacterium spp.* common in zebrafish (*Danio rerio*) research colonies. *J Am Assoc Lab Anim Sci* 56:131–141.
99. Metcalf Pate KA, Rice KA, Wrighten R, Watson J. 2011. Effect of sampling strategy on the detection of fur mites within a naturally infested colony of mice (*Mus musculus*). *J Am Assoc Lab Anim Sci* 50:337–343.
100. Miedel EL, Ragland NH, Engelman RW. 2018. Facility-wide eradication of *Corynebacterium bovis* by using PCR-validated vaporized hydrogen peroxide. *J Am Assoc Lab Anim Sci* 57:465–476. <https://doi.org/10.30802/AALAS-JAALAS-17-000135>.
101. Miller M, Ritter B, Zorn J, Brielmeier M. 2016. Exhaust air dust monitoring is superior to soiled bedding sentinels for detection of *Pasteurella pneumotropica* in individually ventilated cage systems. *J Am Assoc Lab Anim Sci* 55:775–781.
102. Miller M, Brielmeier M. 2017. Environmental samples make soiled bedding sentinels dispensable for hygienic monitoring of IVC-reared mouse colonies. *Lab Anim* 52:233–239. <https://doi.org/10.1177/0023677217739329>.
103. Miller M, Sabrautzi S, Beyerlein A, Brielmeier M. 2019. Combining fish and environmental PCR for diagnostics of diseased laboratory zebrafish in recirculating systems. *PLoS One* 14:1–9. <https://doi.org/10.1371/journal.pone.0222360>.
104. Mocho JP. 2016. Three-dimensional screen: A comprehensive approach to the health monitoring of zebrafish. *Zebrafish* 13 S1:S132–S137. <https://doi.org/10.1089/zeb.2015.1200>.
105. Montonye DR, Ericsson AC, Busi SB, Lutz C, Wardwell K, Franklin CL. 2018. Acclimation and institutionalization of the mouse microbiota following transportation. *Front Microbiol* 9:1–13. <https://doi.org/10.3389/fmicb.2018.01085>.
106. Mullegama SV, Alberti MO, Au C, Li Y, Tou T, Tomasian V, Xian RR. 2019. Nucleic acid extraction from human biological samples. *Methods Mol Biol* 1897:359–383. https://doi.org/10.1007/978-1-4939-8935-5_30.
107. Nashat MA, Luchins KR, Lephed ML, Riedel ER, Izdebska JN, Lipman NS. 2017. Characterization of *Demodex musculi* infestation, associated comorbidities, and topographic distribution in a mouse strain with defective adaptive immunity. *Comp Med* 67:315–329.
108. Nashat MA, Ricart Arbona RJ, Riedel ER, Francino O, Ferrer L, Luchins KR, Lipman NS. 2018. Comparison of diagnostic methods and sampling sites for the detection of *Demodex musculi*. *J Am Assoc Lab Anim Sci* 57:173–185.
109. Ng TFF, Kondov NO, Hayashimoto N, Uchida R, Cha Y, Beyer AI, Wong W, Pesavento PA, Suemizu H, Meunch MO, Delwart EL. 2013. Identification of an astrovirus commonly infecting laboratory mice in the US and Japan. *PLoS One* 8:1–9.
110. Nilsson HO, Ouis IS, Stenram U, Ljungh A, Moran AP, Wadström T, Al-Soud WA. 2004. High prevalence of *Helicobacter* species detected in laboratory mouse strains by multiplex PCR-denaturing gradient gel electrophoresis and pyrosequencing. *J Clin Microbiol* 42:3781–3788. <https://doi.org/10.1128/JCM.42.8.3781-3788.2004>.
111. Ooi JH, Waddell A, Lin Y, Albert I, Rust LT, Holden V, Cantorna MT. 2014. Dominant effects of the diet on the microbiome and the local and systemic immune response in mice. *PLoS One* 9:1–10. <https://doi.org/10.1371/journal.pone.0086366>.
112. Ouellet M, Cowan M, Laporte A, Faubert S, Heon H. 2011. Implementation of a PCR assay of *Pasteurella pneumotropica* to accurately screen for contaminated laboratory mice. *Lab Anim (NY)* 40:305–312. <https://doi.org/10.1038/labani1011-305>.
113. Oz HS, Hughes WT. 1999. DNA amplification of nasopharyngeal aspirates in rats: A procedure to detect *Pneumocystis carinii*. *Microb Pathog* 27:119–121. <https://doi.org/10.1006/mpat.1999.0292>.
114. Parel JDC, Galula JU, Ooi H. 2008. Characterization of rDNA sequences from *Syphacia obvelata*, *Syphacia muris*, and *Aspiculuris tetraptera* and development of a PCR-based method for identification. *Vet Parasitol* 153:379–383. <https://doi.org/10.1016/j.vetpar.2008.02.001>.
115. Parker KD, Albeke SE, Gigley JP, Goldstein AM, Ward NL. 2018. Microbiome composition in both wild-type and disease model mice is heavily influenced by mouse facility. *Front Microbiol* 9:1–13. <https://doi.org/10.3389/fmicb.2018.01598>.
116. Parkinson CM, O'Brien A, Albers TM, Simon MA, Clifford CB, Pritchett-Corning KR. 2011. Diagnosis of ecto- and endoparasites in laboratory rats and mice. *J Vis Exp* 5. <https://doi.org/10.3791/2767>.
117. Pritchett-Corning KR, Prins J, Feinstein R, Goodwin J, Nicklas W, Riley L, Federation of Laboratory Animal Science Associations, American Association for Laboratory Animal Science. 2014. AALAS/FELASA working group on health monitoring of rodents for animal transfer. *J Am Assoc Lab Anim Sci* 53:633–640.
118. Ragland NH, Miedel EL, Engelman RW. 2019. PCR prevalence of murine opportunistic microbes and their mitigation by using vaporized hydrogen peroxide. *J Am Assoc Lab Anim Sci* 58:208–215. <https://doi.org/10.30802/AALAS-JAALAS-18-000112>.
119. Rausch P, Basic M, Batra A, Bischoff SA, Blaut M, Clavel T, Glasner J, Gopalakrishnan S, Grassl GA, Gunther C, Haller D, Hirose M, Ibrahim S, Loh G, Mattner J, Nagel S, Pabst O, Schmidt F, Siegmund B, Strowig T, Volynets V, Wirtz S, Zeissig S, Zeissig Y, Bleich A, Baines JF. 2016. Analysis of factors contributing to variation in the C57BL/6J fecal microbiota across German animal facilities. *Int J Med Microbiol* 306:343–355. <https://doi.org/10.1016/j.ijmm.2016.03.004>.
120. Regan RD, Fenyk-Melody JE, Tran SM, Chen G, Stocking KL. 2016. Comparison of submental blood collection with the

- retroorbital and submandibular methods in mice (*Mus musculus*). *J Am Assoc Lab Anim Sci* 55:570–576.
121. **Rice KA, Albacarys LK, Metcalf Pate KA, Perkins C, Henderson KS, Watson J.** 2013. Evaluation of diagnostic methods for *Myocoptes musculinus* according to age and treatment status of mice (*Mus musculus*). *J Am Assoc Lab Anim Sci* 52:773–781.
122. **Riley LK, Franklin CL, Hook RR Jr, Besch-Williford C.** 1996. Identification of murine *Helicobacters* by PCR and restriction enzyme analyses. *J Clin Microbiol* 34:942–946. <https://doi.org/10.1128/JCM.34.4.942-946.1996>.
123. **Rowe WP, Hartley JW, Estes JD, Huebner RJ.** 1959. Studies of mouse polyoma virus infection. 1. procedures for quantitation and detection of virus. *J Exp Med* 109:379–391. <https://doi.org/10.1084/jem.109.4.379>.
124. **Sagaidak S, Taibi A, Wen B, Comelli EM.** 2016. Development of a real-time PCR assay for quantification of *Citrobacter rodentium*. *J Microbiol Methods* 126:76–77. <https://doi.org/10.1016/j.mimet.2016.05.008>.
125. **Sanchez S, Tyler K, Rozengurt N, Lida J.** 1994. Comparison of a PCR-based diagnostic assay for *Mycoplasma pulmonis* with traditional detection techniques. *Lab Anim* 28:249–256. <https://doi.org/10.1258/002367794780681570>.
126. **Sanders JL, Kent ML.** 2011. Development of a sensitive assay for the detection of *Pseudoloma neurophilia* in laboratory populations of the zebrafish *Danio rerio*. *Dis Aquat Organ* 96:145–156. <https://doi.org/10.3354/dao02375>.
127. **Scott ME, Gibbs HC.** 1986. Long-term population dynamics of pinworms (*Syphacia obvelata* and *Aspiculuris tetraptera*) in mice. *J Parasitol* 72:652–662. <https://doi.org/10.2307/3281452>.
128. **Smith AL.** 1986. Methods for potential application to rodent virus isolation and identification, p 753–776. Chapter 35. In: Bhatt PN, Jacoby RO, Morse HC, New AE, editors. *Viral and mycoplasmal infections of laboratory rodents: effects on biomedical research*. Orlando (FL): Academic Press.
129. **Smith AL.** 1986. Serologic tests for detection of antibody to rodent viruses, p 731–751. Chapter 34. In: Bhatt PN, Jacoby RO, Morse HC, New AE, editors. *Viral and mycoplasmal infections of laboratory rodents: effects on biomedical research*. Orlando (FL): Academic Press.
130. **Smith PC, Zeiss CJ, Beck AP, Scholz JA.** 2016. *Demodex musculi* infestation in genetically immunomodulated mice. *Comp Med* 66:278–285.
131. **Sofi MH, Gudi R, Karumuthil-Melethil S, Perez N, Johnson BM, Vasu C.** 2014. pH of drinking water influences the composition of gut microbiome and type 1 diabetes incidence. *Diabetes* 63:632–644. <https://doi.org/10.2337/db13-0981>.
132. **Thoene-Reineke C, Fischer A, Friese C, Briesemeister D, Gobel UB, Kammertoens T, Bereswill S, Heimesaat MM.** 2014. Composition of intestinal microbiota in immune-deficient mice kept in three different housing conditions. *PLoS One* 9:1–8. <https://doi.org/10.1371/journal.pone.0113406>.
133. **Wagner AM, Loganbill JK, Besselsen DG.** 2004. Detection of lactate dehydrogenase-elevating virus by use of a fluorogenic nuclease reverse transcriptase-polymerase chain reaction assay. *Comp Med* 54:288–292.
134. **Ward JM, Fox JG, Anver MR, Haines DC, George CV, Collins MJ Jr, Gorelick PL, Nagashima K, Gonda MA, Gilden RV.** 1994. Chronic active hepatitis and associated liver tumors in mice caused by a persistent bacterial infection with a novel *Helicobacter* species. *J Natl Cancer Inst* 86:1222–1227. <https://doi.org/10.1093/jnci/86.16.1222>.
135. **Weiss EE, Evans KD, Griffey SM.** 2012. Comparison of a fur mite PCR assay and the tape test for initial and posttreatment diagnosis during a natural infection. *J Am Assoc Lab Anim Sci* 51:574–578.
136. **Whary MT, Fox JG.** 2006. Detection, eradication, and research implications of *Helicobacter* infections in laboratory rodents. *Lab Anim (NY)* 35:25–27, 30–36. <https://doi.org/10.1038/lab-an0706-25>.
137. **Zeiss C, Neaderland M, Yang FC, Terwilliger G, Compton S.** 2012. Fungal polymerase chain reaction testing in equine ulcerative keratitis. *Vet Ophthalmol* 16:341–351. <https://doi.org/10.1111/vop.12004>.
138. **Zorn J, Ritter B, Miller M, Kraus M, Northrup E, Briemeier M.** 2016. Murine norovirus detection in the exhaust air of IVCs is more sensitive than serological analysis of soiled bedding sentinels. *Lab Anim* 51:301–310. <https://doi.org/10.1177/0023677216661586>.